

## DOCUMENT RESUME

ED 465 945

CG 031 845

AUTHOR Simpson, Chris  
TITLE Self-Mutilation. ERIC/CASS Digest.  
INSTITUTION ERIC Clearinghouse on Counseling and Student Services,  
Greensboro, NC.  
SPONS AGENCY Office of Educational Research and Improvement (ED),  
Washington, DC.  
REPORT NO EDO-CG-01-10  
PUB DATE 2001-12-00  
NOTE 4p.  
CONTRACT ED-99-CO-0014  
AVAILABLE FROM ERIC Counseling and Student Services Clearinghouse,  
University of North Carolina at Greensboro, 201 Ferguson  
Building, P.O. Box 26171, Greensboro, NC 27402-6171: Tel:  
336-334-4114; Tel: 800-414-9769 (Toll Free); Fax:  
336-334-4116; e-mail: ericcass@uncg.edu. For full text:  
<http://ericcass.uncg.edu>.  
PUB TYPE ERIC Publications (071) -- ERIC Digests in Full Text (073)  
EDRS PRICE MF01/PC01 Plus Postage.  
DESCRIPTORS Behavior Disorders; \*Client Characteristics (Human  
Services); \*Clinical Diagnosis; Counseling Techniques;  
Emotional Disturbances; \*Mental Health Workers; \*Self  
Injurious Behavior  
IDENTIFIERS ERIC Digests

## ABSTRACT

Self-mutilation has been most commonly seen as a diagnostic indicator for borderline personality disorder. However, practitioners have more recently observed self-harming behavior among those individuals diagnosed with bipolar disorder, obsessive-compulsive disorder, eating disorders, multiple personality disorder, borderline personality disorder, schizophrenia, and most recently, with adolescents and young adults. The increased observance of these behaviors has left many mental health professionals calling for self-mutilation to have its own diagnosis in the Diagnostic and Statistical Manual of Mental Disorders. The phenomenon is often difficult to define and easily misunderstood. This digest presents a definition of self-mutilation, reasons for self-mutilating behavior, characteristics of individuals who self-mutilate, common misconceptions of self-mutilation, and treatment options. It is noted in conclusion that although therapeutic interventions have improved substantially over the past two decades, further study is imperative to insure that those who practice the behavior continue to receive effective care. (GCP)



# *Self-Mutilation*

## *ERIC Digest*

By

Chris Simpson

December 2001

EDO-CG-01-10

ERIC Clearinghouse for Counseling and Student Services

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC)

- ☐ This document has been reproduced as received from the person or organization originating it.
- ☐ Minor changes have been made to improve reproduction quality.

- 
- Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

# Self-Mutilation

Chris Simpson

December, 2001  
EDO-CG-01-10

## Introduction

Suyemoto and MacDonald (1995) reported that the incidence of self-mutilation occurred in adolescents and young adults between the ages of 15 and 35 at an estimated 1,800 individuals out of 100,000. The incidence among inpatient adolescents was an estimated 40%. Self-mutilation has been most commonly seen as a diagnostic indicator for Borderline Personality Disorder, a characteristic of Stereotypic Movement Disorder (associated with autism and mental retardation) and attributed to Factitious Disorders. However, practitioners have more recently observed self-harming behavior among those individuals diagnosed with bipolar disorder, obsessive-compulsive disorder, eating disorders, multiple personality disorder, borderline personality disorder, schizophrenia, and most recently, with adolescents and young adults. The increased observance of these behaviors has left many mental health professionals calling for self-mutilation to have its own diagnosis in the *Diagnostic and Statistical Manual of Mental Disorders* (Zila & Kiselica, 2001). The phenomenon is often difficult to define and easily misunderstood.

## Definition of Self-Mutilation

Several definitions of this phenomenon exist. In fact, researchers and mental health professionals have not agreed upon one term to identify the behavior. Self-harm, self-injury, and self-mutilation are often used interchangeably.

Some researchers have categorized self-mutilation as a form of self-injury. Self-injury is characterized as any sort of self-harm that involves inflicting injury or pain on one's own body. In addition to self-mutilation, examples of self-injury include: hair pulling, picking the skin, excessive or dangerous use of mind-altering substances such as alcohol, and eating disorders.

Favazza and Rosenthal (1993) identify pathological self-mutilation as the deliberate alteration or destruction of body tissue without conscious suicidal intent. A common example of self-mutilating behavior is cutting the skin with a knife or razor until pain is felt or blood has been drawn. Burning the skin with an iron, or more commonly with the ignited end of a cigarette, is also a form of self-mutilation.

Self-mutilating behavior does exist within a variety of populations. For the purpose of accurate identification, three different types of self-mutilation have been identified:

1) superficial or moderate; 2) stereotypic; and 3) major. Superficial or moderate self-mutilation is seen in individuals diagnosed with personality disorders (i.e. borderline personality disorder). Stereotypic self-mutilation is often associated with mentally delayed individuals. Major self-mutilation, more rarely documented than the two previously mentioned categories, involves the amputation of the limbs or genitals. This category is most commonly associated with pathology (Favazza & Rosenthal,

1993). The remaining portion of this digest will focus on superficial or moderate self-mutilation.

Additionally, self-injurious behavior may be divided into two dimensions: nondissociative and dissociative. Self-mutilative behavior often stems from events that occur in the first six years of a child's development.

Nondissociative self-mutilators usually experience a childhood in which they are required to provide nurturance and support for parents or caretakers. If a child experiences this reversal of dependence during formative years, that child perceives that she can only feel anger toward self, but never toward others. This child experiences rage, but cannot express that rage toward anyone but him or herself. Consequently, self-mutilation will later be used as a means to express anger.

Dissociative self-mutilation occurs when a child feels a lack of warmth or caring, or cruelty by parents or caretakers. A child in this situation feels disconnected in his/her relationships with parents and significant others. Disconnection leads to a sense of "mental disintegration." In this case, self-mutilative behavior serves to center the person (Levenkron, 1998, p. 48).

## Reasons for Self-Mutilating Behavior

Individuals who self-injure often have suffered sexual, emotional, or physical abuse from someone with whom a significant connection has been established such as a parent or sibling. This often results in the literal or symbolic loss or disruption of the relationship. The behavior of superficial self-mutilation has been described as an attempt to escape from intolerable or painful feelings relating to the trauma of abuse.

The person who self-harms often has difficulty experiencing feelings of anxiety, anger, or sadness. Consequently, cutting or disfiguring the skin serves as a coping mechanism. The injury is intended to assist the individual in dissociating from immediate tension (Stanley, Gameroff, Michaelson & Mann, 2001).

## Characteristics of Individuals Who Self-Mutilate

Self-mutilating behavior has been studied in a variety of racial, chronological, ethnic, gender, and socioeconomic populations. However, the phenomenon appears most commonly associated with middle to upper class adolescent girls or young women.

People who participate in self-injurious behavior are usually likeable, intelligent, and functional. At times of high stress, these individuals often report an inability to think, the presence of unexpressable rage, and a sense of powerlessness. An additional characteristic identified by researchers and therapists is the inability to verbally express feelings.

Some behaviors found in other populations have been mistaken for self-mutilation. Individuals who have tattoos or piercings are often falsely accused of being self-mutilators. Although these practices have varying degrees of social acceptability, the behavior is not typical of self-mutilation. The majority of these persons tolerate pain for the purpose of attaining a finished product like a piercing or tattoo. This differs from the individual who self-mutilates for whom pain experienced from cutting or damaging the skin is sought as an escape from intolerable affect (Levenkron, 1998).

### Common Misconceptions of Self-Mutilation

#### *Suicide*

Stanley et al., (2001) report that approximately 55%-85% of self-mutilators have made at least one attempt at suicide. Although suicide and self-mutilation appear to possess the same intended goal of pain relief, the respective desired outcomes of each of these behaviors is not entirely similar.

Those who cut or injure themselves seek to escape from intense affect or achieve some level of focus. For most members of this population, the sight of blood and intensity of pain from a superficial wound accomplish the desired effect, dissociation or management of affect. Following the act of cutting, these individuals usually report feeling better (Levenkron, 1998).

Motivation for committing suicide is not usually characterized in this manner. Feelings of hopelessness, despair, and depression predominate. For these individuals, death is the intent. Consequently, though the two behaviors possess similarities, suicidal ideation and self-mutilation may be considered distinctly different in intent.

#### *Attention-seeking behavior*

Levenkron (1998) reports that individuals who self-mutilate are often accused of "trying to gain attention." Although self-mutilation may be considered a means of communicating feelings, cutting and other self-harming behavior tends to be committed in privacy. In addition, self-harming individuals will often conceal their wounds. Revealing self-inflicted injuries will often encourage other individuals to attempt to stop the behavior. Since cutting serves to dissociate the individual from feelings, drawing attention to wounds is not typically desired. Those individuals who commit self-harm with the intent of gaining attention are conceptualized differently from those who self-mutilate.

#### *Dangerousness to others*

Another reported misconception is that those individuals who commit self-harm are a danger to others. Although self-mutilation has been identified as a characteristic of individuals suffering from a variety of diagnosed pathology, most of these individuals are functional and pose no threat to the safety of other persons.

### Treatment of the Individual Who Self-Mutilates

Methods employed to treat those persons who self-mutilate range on a continuum from successful to ineffective. Those treatment methods that have shown effectiveness in working with this population include: art therapy, activity therapy, individual counseling, and support groups. An important skill of the professional working with a self-harming individual is the ability to look at wounds without grimacing or passing judgment (Levenkron, 1998). A setting that promotes the healthy expression of emotions, and counselor patience and willingness to examine wounds is the common bond among these progressive interventions (Levenkron, 1998; Zila & Kiselica, 2001).

### Conclusion

Research shows that self-mutilation has been in existence far longer than the understanding and accurate conceptualization of the phenomenon. Therapeutic interventions have improved substantially over the past two decades. However, further study is imperative to insure that those who practice the behavior continue to receive effective care.

### References

- Favaro, A. & Santonastaso, P. (2000). Self-injurious behavior in anorexia nervosa. *The Journal of Nervous and Mental Disease*, 188(8), 537-542.
- Favazza, A.R. & Rosenthal, R. J. (1993). Diagnostic issues in self-mutilation. *Hospital and Community Psychiatry*, 44, 134-140.
- Levenkron, S. (1998). *Cutting*. New York, NY: W. W. Norton and Company.
- Stanley, B., Gameroff, M. J., Michalsen, V., & Mann, J. J. (2001). Are suicide attempters who self-mutilate a unique population? *American Journal of Psychiatry*, 158(3), 427-432.
- Suyemoto, K. L. & MacDonald, M. L. (1995). Self-cutting in female adolescents. *Psychotherapy*, 32(1), 162-171.
- Zila, L. M. & Kiselica, M. S. (2001). Understanding and counseling self-mutilation in female adolescents and young adults. *Journal of Counseling & Development*, 79, 46-52.

*Chris Simpson is an assistant professor at San Francisco State University, San Francisco, California.*

ERIC Digests are in the public domain and may be freely reproduced and disseminated. This publication was funded by the U.S. Department of Education, Office of Educational Research and Improvement, Contract No. ED-99-CO-0014. Opinions expressed in this report do not necessarily reflect the positions of the U.S. Department of Education, OERI, or ERIC/CASS.

For information on other ERIC/CASS products and services, please call toll-free (800) 414-9769 or (336) 334-4114 or fax (336) 334-4116 or write ERIC/CASS, School of Education, University of North Carolina at Greensboro, Greensboro, NC 27402.



**U.S. Department of Education**  
*Office of Educational Research and Improvement (OERI)*  
*National Library of Education (NLE)*  
*Educational Resources Information Center (ERIC)*



## **NOTICE**

### **Reproduction Basis**



This document is covered by a signed "Reproduction Release (Blanket)" form (on file within the ERIC system), encompassing all or classes of documents from its source organization and, therefore, does not require a "Specific Document" Release form.



This document is Federally-funded, or carries its own permission to reproduce, or is otherwise in the public domain and, therefore, may be reproduced by ERIC without a signed Reproduction Release form (either "Specific Document" or "Blanket").